

UK-1, A NOVEL CYTOTOXIC METABOLITE FROM *Streptomyces* sp. 517-02

## II. STRUCTURAL ELUCIDATION

KOZO SHIBATA, MASARU KASHIWADA, MASASHI UEKI and MAKOTO TANIGUCHI\*

Faculty of Science, Osaka City University,  
3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558, Japan

(Received for publication January 21, 1993)

The structure of UK-1 isolated from the mycelium of *Streptomyces* sp. 517-02 was elucidated to be a novel benzoxazole dimer derivative (1) on the basis of spectroscopic methods.

A novel metabolite with potent cytotoxic activity against B16, HeLa and P388 cells, UK-1, was isolated from the mycelium of *Streptomyces* sp. 517-02 as described in a previous paper<sup>1)</sup>. The structure of UK-1 was elucidated to be a dimeric benzoxazole derivative constituted of two moles of 3-hydroxyanthranilic acid and one mole of salicylic acid on the basis of some spectroscopic methods (Fig. 1). The structure determination studies of UK-1 are described in this paper.

## Results and Discussion

The IR spectrum of UK-1 (1) described in a previous paper showed a strong absorption based on ester group at  $1725\text{ cm}^{-1}$  and the UV spectrum of 1 suggested the existence of conjugated system in the molecule. The molecular formula of 1 was determined as  $\text{C}_{22}\text{H}_{14}\text{N}_2\text{O}_5$  from the HREI-MS ( $\text{M}^+$ :  $m/z$  386.0913, Calcd for  $\text{C}_{22}\text{H}_{14}\text{N}_2\text{O}_5$ , 386.0903) as the base peak and  $^{13}\text{C}$  NMR spectral data. Other fragment ions were observed at  $m/z$  354.0606 (Calcd for  $\text{C}_{22}\text{H}_{14}\text{N}_2\text{O}_5 \cdot \text{CH}_3\text{OH}$ , 354.0572) and  $m/z$  328.0888 (Calcd for  $\text{C}_{22}\text{H}_{14}\text{N}_2\text{O}_5 \cdot \text{HCOOCH}_3$ , 328.0928) in the HREI-MS spectrum of 1. The absorption of a hydroxyl group could not be observed in the IR spectrum of 1 but the signal based on a strong hydrogen bonded hydroxyl group appeared at  $\delta$  11.9 ppm in the  $^1\text{H}$  NMR spectrum in  $\text{CDCl}_3$ . Moreover, 1 afforded its mono-methyl ether, Me-UK-1 (2), by methylation with methyl iodide and anhydrous potassium carbonate in dry acetone. The IR spectrum of 2 showed the absorption based on an ester group at  $\nu_{\text{max}}$   $1710\text{ cm}^{-1}$ . In the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of 2, the signals of a methoxyl group appeared at  $\delta$  4.09 and  $\delta$  56.22, respectively showing to be the monomethyl derivative of 1. Alkaline hydrolysis of 1 furnished the corresponding carboxylic acid, DeMe-UK-1 (3). The absorption based on a carboxyl group appeared at  $\nu_{\text{max}}$   $2500\sim 3100$  and  $1690\text{ cm}^{-1}$  in the IR spectrum of 3 and the signal of a methoxycarbonyl group disappeared in the  $^1\text{H}$  NMR.

The  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectral data of 1, 2 and 3 are shown in Table 1. The assignments of proton and carbon signals were done using  $^1\text{H}$ - $^1\text{H}$  COSY and  $^1\text{H}$ - $^{13}\text{C}$  COSY spectra (Fig. 2) and

Fig. 1. Structures of UK-1 (1) and its derivatives.




Table 1. NMR spectral data for benzoxazoles.

No.	UK-1 (1) <sup>a</sup>		Me-UK-1 (2) <sup>a</sup>		DeMe-UK-1 (3) <sup>b</sup>		2-o-Methoxybenzoxazole (5) <sup>a</sup>		2-o-Hydroxybenzoxazole (6) <sup>a</sup>	
	<sup>13</sup> C shift	<sup>1</sup> H shift	<sup>13</sup> C shift	<sup>1</sup> H shift	<sup>13</sup> C shift	<sup>1</sup> H shift	<sup>13</sup> C shift	<sup>1</sup> H shift	<sup>13</sup> C shift	<sup>1</sup> H shift
1	151.41		151.86		151.83					
2	141.59		141.64		142.05					
3	122.67		122.39		122.98					
4	127.41	8.07 dd	127.11	8.06 dd	127.82	8.39 d				
5	124.83	7.44 t	124.47	7.44 t	125.32	7.47 t				
6	114.96	7.82 dd	115.34	7.91 dd	114.71	7.85 d				
7	161.81		163.20		159.93 <sup>c</sup>					
8	117.77		118.45		118.00		120.25	7.59 d	119.55	7.69 d
9	138.74		140.75		138.78		142.24		140.14	
10	149.98		151.47		150.28		150.53		149.23	
11	113.69	7.72 dd	113.88	7.78 dd	114.20	7.79 d	110.48	7.81 d	110.64	7.71 d
12	125.22	7.48 t	124.76	7.50 t	125.69	7.47 t	124.28	7.35 t	125.01	7.36 t
13	125.33	8.31 dd	125.79	8.43 dd	125.88	8.42 d	124.90	7.32 t	125.37	7.33 t
14	164.62		163.89		164.57		161.87		162.98	
15	110.06		116.00		110.57		116.50		110.64	
16	159.68		159.03		159.93 <sup>c</sup>		158.62		158.85	
17	117.84	7.15 d	112.34	7.10 d	118.16	7.31 d	112.26	7.08 d	117.47	7.12 d
18	134.24	7.46 t	133.37	7.54 t	134.71	7.51 t	132.73	7.49 t	133.56	7.42 t
19	119.56	7.00 td	120.83	7.13 t	120.04	7.05 t	120.77	7.07 t	119.27	6.98 t
20	127.21	8.01 dd	131.95	8.32 d	127.65	8.04 d	131.40	8.13 d	127.15	8.01 d
C-O	166.12		165.86		167.81					
OMe	52.53	4.17 s	52.32	4.10 s			56.26	4.01 s		
OMe			56.22	4.09 s						
OH		11.90 s								11.42 s

<sup>a</sup> In CDCl<sub>3</sub>.<sup>b</sup> In pyridine-d<sub>5</sub>.<sup>c</sup> Overlapped signal.



Fig. 2.  $^1\text{H}$ - $^{13}\text{C}$  COSY spectrum of Me-UK-1 (2).

Fig. 3. COLOC spectrum of Me-UK-1 (2).




correlation spectroscopy *via* long-range couplings (COLOC) measurement (Fig. 3). These results revealed the partial structure A, B and C in 2 as shown in Fig. 4, and some possible formulae as the structure of 1 were estimated by the combination of these partial structures.


The  $^{13}\text{C}$  NMR parameters in the benzene ring of 1 are consistent with the values calculated on the basis of the chemical shift of carbons in the benzene ring of benzoxazole (4)<sup>2</sup>. Moreover, the benzoxazole derivatives (5 and 6) were prepared from *o*-aminophenol and *o*-anisaldehyde or salicylaldehyde, respectively (Scheme 1)<sup>3</sup>, and the  $^{13}\text{C}$  chemical shifts of these derivatives were in good accordance with those of 1 and 2 (Table 1).

From these results, the structure of UK-1 was deduced to be formula 1, the novel benzoxazole dimer derivative.


Fig. 4. Partial structures of Me-UK-1 (2) and the correlation of  $^1\text{H}$ - $^{13}\text{C}$  long-range couplings.



Scheme 1.



Scheme 2. Estimation of biosynthesis for UK-1.



Both of the fragments A and B of 1 are related to 3-hydroxyanthranilic acid, one of the catabolic products *via* kynurenine and the fragment C is a reduced product of salicylic acid. The benzoxazoles 5 and 6 were easily prepared by oxidation of SCHIFF's bases derived from *o*-aminophenol and corresponding aldehydes. It seems that UK-1 was biosynthesized by oxidation of the SCHIFF's base prepared from methyl 3-hydroxyanthranilate and 3-hydroxyanthranilaldehyde produced in the decomposition pathway of L-tryptophan, followed to the preparation of SCHIFF's base with salicylaldehyde and the oxidative ring closure reaction of SCHIFF's base (Scheme 2). Studies on the biosynthesis of 1 are now in progress.

### Experimental

#### MS and NMR

EI-MS and HREI-EI-MS spectra were obtained with a JEOL-JMS-AX 500 mass spectrometer. All NMR spectra were recorded on a JEOL-JNM-GX-400 spectrometer operating at 400 MHz for  $^1\text{H}$  NMR and 100 MHz for  $^{13}\text{C}$  NMR. Tetramethylsilane was used as an internal reference for  $^1\text{H}$  NMR in the  $\text{CDCl}_3$  solution. For the  $^{13}\text{C}$  chemical shift reference, the  $^{13}\text{C}$  peak at  $\delta$  77.03 ppm of  $\text{CDCl}_3$  was used. The  $^1\text{H}$  peak at  $\delta$  7.0 ppm and  $^{13}\text{C}$  peak at  $\delta$  122.4 ppm of pyridine- $d_5$  were used as the internal references for NMR measurement in pyridine- $d_5$ .

#### Methylation of 1

Anhydrous potassium carbonate (2 g) was suspended in a solution of 50 mg of 1 and 0.5 ml of methyl iodide in 20 ml of dry acetone, and the mixture was refluxed for 5 hours. After filtration, acetone was evaporated and the residual mixture was dissolved in  $\text{CH}_2\text{Cl}_2$ , followed to chromatographical purification on a silica gel column. By recrystallization with methanol, 2 was obtained as colorless needles in quantitative yield. 2; MP 145~147°C, IR (nujol) 1710, 1600, 1590, 1580, 1540  $\text{cm}^{-1}$ , HREI-MS ( $\text{M}^+$ ) =  $m/z$  400.1067 (Calcd for  $\text{C}_{23}\text{H}_{18}\text{N}_2\text{O}_5$ , 400.1075), NMR, see Table 1.

#### Alkaline Hydrolysis of 1

Alkaline hydrolysis of 1 with aqueous NaOH in a pyridine solution at room temperature afforded 3. 3; MP > 300°C, IR (nujol) 2500~3100, 1690, 1600, 1560  $\text{cm}^{-1}$ , NMR, see Table 1.

#### Preparation of 2-Phenylbenzoxazoles

2-Phenylbenzoxazoles were prepared according to the procedure of STEPHENS and BOWER<sup>3)</sup>. Namely, *o*-aminophenol dissolved in ethanol was mixed with the corresponding aldehyde, boiled for 10 minutes and cooled. The product obtained by filtration and recrystallization from ethanol gave a red SCHIFF's base. The treatment of the SCHIFF's base with lead tetraacetate in glacial acetic acid afforded the 2-phenylbenzoxazole, 4 or 5.

4; The yield of SCHIFF's base from *o*-aminophenol (1.1 g) and *o*-anisaldehyde (1.4 g) was 1.04 g (70%). The dehydrogenation of SCHIFF's base (140 mg) with lead tetraacetate (211 mg) in acetic acid (3.5 ml) afforded 2-*o*-methoxyphenylbenzoxazole (38 mg).  $^1\text{H}$  and  $^{13}\text{C}$  NMR, see Table 1.

5; The yield of SCHIFF's base from *o*-aminophenol (1.1 g) and salicylaldehyde (1.2 g) was 2.24 g (100%). The dehydrogenation of SCHIFF's base (1.23 g) with lead tetraacetate (2.1 g) in acetic acid (20 ml) afforded 2-*o*-hydroxyphenylbenzoxazole.  $^1\text{H}$  and  $^{13}\text{C}$  NMR, see Table 1.

### Acknowledgments

We are grateful to Prof. S. OI of our laboratory for their instructive advice and helpful discussion. We also wish to thank Mr. J. GOUDA and Mr. T. SHIMADA for their kind measurement of elemental analysis and MS experiments.

### References

- 1) UEKI, M.; K. UENO, S. MIYADOH, K. ABE, K. SHIBATA, M. TANIGUCHI & S. OI: UK-1, a novel cytotoxic metabolite

- from *Streptomyces* sp. 517-02. I. Taxonomy, fermentation, isolation, physico-chemical and biological properties. *J. Antibiotics* 46: 1089~1094, 1993
- 2) KALINOWSKI, H.; S. BERGER & S. BRAUN (Eds.): Carbon-13 NMR Spectroscopy. pp. 313~315 and 394, John Wiley & Sons Ltd., 1988
  - 3) STEPHENS, F. F. & J. D. BOWER: The preparation of benzimidazoles and benzoxazoles from SCHIFF's bases. Part I. *J. Chem. Soc.* 1949: 2971~2972, 1949